

CONSTRUCTION OF A PARAMETRIC FAMILY OF DIOPHANTINE TRIPLES IN INTEGERS

Manju Somanath¹, <manjuajil@yahoo.com>
J. Kannan^{2*} <jayram.kannan@gmail.com>
K. Raja³ <rajakonline@gmail.com>

^{1,2,3}Department of Mathematics, National College, 620 001, Tamil Nadu, India

Abstract – Let n be a non-zero integer. A set $\{a_1, a_2, \dots, a_m\}$ of m distinct positive integers is called a Diophantine m – tuples with the property $D(n)$, if $a_i a_j + n$ is a perfect square for all $1 \leq i < j \leq m$. In this paper, we give some sets of polynomial with integer coefficients, such that the product of any two of them added with a quadratic polynomial in $Z(n)$, is a square of a polynomial with integer coefficients.

Keywords – Diophantine triple, Perfect square, Quadratic Polynomial

1 INTRODUCTION

In this paper, we study Diophantine sets, defined as consisting of linear polynomials with the property that, adding the quadratic polynomials $\sigma^2 + 10\sigma + 6$ or $10\omega^2 + 26\omega + 13$, (here $\sigma, \omega \in \mathbf{N}$) to the product of any two of them, one gets a perfect square.

The idea of extending a Diophantine set by joining an integer that preserves the defining property was employed since Euler. Obtaining Dio-pairs is easy: for any integer $r \geq 2$, find a factor a of $r^2 - 1$ and consider it along with the cofactor $\frac{r^2 - 1}{a}$. It requires an extensive work to find all Diophantine triples extending a fixed pair (a, b) . The problem involves solving of Pellian equation for finding a "c" such that ac and bc also preserves the defined property. The case of extending a triple to quadruple have been studied by many mathematicians. In 1993, Dujella proved that if an integer n does not have the form $n = 4k + 2$ and $n \notin S = \{-4, -3, -1, 3, 5, 8, 12, 20\}$, then there exists at least one Diophantine quadruple with property $D(n)$.

2 PRELIMINARY

Many parametric families of Diophantine triples and 4 – tuples are known. for example.

$$\{(a, b, c) : ab + 1 = q^2, c = a + b + 2q\}. \quad (1)$$

** Edited by A. D. Godase (Area Editor).

* Corresponding Author J. Kannan

is a family of Diophantine triples for $0 < a < b, q > 0$, and

$$\{(a, b, c, d) : ab + 1 = x^2, bc + 1 = y^2, ca + 1 = z^2, d = a + b + c + 2abc + 2xyz\} \quad (2)$$

is a family of Diophantine 4 - *tuples*. Note that (1) shows how to pass from a Diophantine pair (a, b) to a Diophantine triple (a, b, c) , and (2) shows how to pass from a Diophantine triple (a, b, c) to a Diophantine 4 - tuple (a, b, c, d) .

Let us briefly consider the problem of extending a Diophantine pair (a, b) , where $ab + 1 = q^2$, to a Diophantine triple (a, b, c) . We shall assume that a and b are coprime, and we require an integer c such that $ac + 1 = X^2$ and $bc + 1 = Y^2$ for some integer X and Y . If such a, c exist, then X and Y must be solutions of the Diophantine equation $bX^2 - aY^2 = b - a$. Conversely, if X and Y are solutions of this equation then a divides $X^2 - 1$ (because a and b are coprime) and so we can define c by $ac + 1 = X^2$ and $bc + 1 = Y^2$ and (a, b, c) is a Diophantine triple. As the Diophantine equation behave erratically with respect to the coefficients, it therefore seems unlikely that we can parametrise all Diophantine triples of the form (a, b, x) . In this argument, the Diophantine triples, in (1) correspond to the solutions $X = a + q$ and $Y = b + q$.

3 DISCUSSION AND RESULTS

Construction of a Diophantine triple (α, β, γ) of linear polynomials such that the product of any two of them added with quadratic polynomials explained in choices 1 and 2, leaves a perfect square.

3.1 Choice 1

For the different pairs of linear polynomials, we investigate for its extendability as a triple satisfying the property $D(\sigma^2 + 10\sigma + 6)$.

3.1.1 Case 1

Consider the $\{\alpha, \beta\}$ where $\alpha = 3\sigma + 1$ and $\beta = 5\sigma + 3$ satisfying $D(\sigma^2 + 10\sigma + 6)$. To search for its extendability, let γ be any other polynomial with the same property. This can be written as $\alpha\gamma + \sigma^2 + 10\sigma + 6 = \rho^2$ and $\beta\gamma + \sigma^2 + 10\sigma + 6 = \eta^2$. Applying the linear transformation, $\rho = \mu + (3\sigma + 1)\nu$ and $\eta = \mu + (5\sigma + 3)\nu$ and eliminating γ , we get the Pellian equation $\mu^2 - (15\sigma^2 + 14\sigma + 3)\nu^2 = \sigma^2 + 10\sigma + 6$, with initial solution $\mu_0 = 4\sigma + 3, \nu_0 = 1$. The values of μ_0 and ν_0 can be employed to find the polynomial γ as $\gamma = 16\sigma + 10$, satisfying the property $D(\sigma^2 + 10\sigma + 6)$.

3.1.2 Case 2

Consider the $\{\beta, \gamma\}$ where $\beta = 5\sigma + 3$ and $\gamma = 16\sigma + 10$ satisfying $D(\sigma^2 + 10\sigma + 6)$. To search for its extendability, let δ be any other polynomial with the same property. This can be written as $\beta\delta + \sigma^2 + 10\sigma + 6 = \rho^2$ and $\gamma\delta + \sigma^2 + 10\sigma + 6 = \eta^2$. Applying the linear transformation, $\rho = \mu + (5\sigma + 3)\nu$ and $\eta = \mu + (16\sigma + 10)\nu$ and eliminating δ , we get the Pellian equation $\mu^2 - (80\sigma^2 + 98\sigma + 30)\nu^2 = \sigma^2 + 10\sigma + 6$, with initial solution $\mu_0 = 9\sigma + 6, \nu_0 = 1$. The values of μ_0 and ν_0 can be employed to find the polynomial δ as $\delta = 39\sigma + 25$, satisfying the property $D(\sigma^2 + 10\sigma + 6)$.

3.1.3 Case 3

Consider the $\{\gamma, \delta\}$ where $\gamma = 16\sigma + 10$ and $\delta = 39\sigma + 25$ satisfying $D(\sigma^2 + 10\sigma + 6)$. To search for its extendability, let ξ be any other polynomial with the same property. This can be written as $\gamma\xi + \sigma^2 + 10\sigma + 6 = \rho^2$ and $\delta\xi + \sigma^2 + 10\sigma + 6 = \eta^2$. Applying the linear transformation, $\rho = \mu + (16\sigma + 10)\nu$ and $\eta = \mu + (39\sigma + 25)\nu$ and eliminating ξ , we get the Pellian equation $\mu^2 - (624\sigma^2 + 790\sigma + 250)\nu^2 = \sigma^2 + 10\sigma + 6$, with initial solution $\mu_0 = 25\sigma + 16, \nu_0 = 1$. The values of μ_0 and ν_0 can be employed to find the polynomial ξ as $\xi = 105\sigma + 67$, satisfying the property $D(\sigma^2 + 10\sigma + 6)$.

3.1.4 Case 4

Consider the $\{\delta, \xi\}$ where $\delta = 39\sigma + 25$ and $\xi = 105\sigma + 67$ satisfying $D(\sigma^2 + 10\sigma + 6)$. To search for its extendability, let ζ be any other polynomial with the same property. This can be written as $\delta\zeta + \sigma^2 + 10\sigma + 6 = \rho^2$ and $\xi\zeta + \sigma^2 + 10\sigma + 6 = \eta^2$.

Applying the linear transformation, $\rho = \mu + (39\sigma + 25)\nu$ and $\eta = \mu + (105\sigma + 67)\nu$ and eliminating ζ , we get the Pellian equation $\mu^2 - (4095\sigma^2 + 5238\sigma + 1675)\nu^2 = \sigma^2 + 10\sigma + 6$, with initial solution $\mu_0 = 64\sigma + 41, \nu_0 = 1$. The values of μ_0 and ν_0 can be employed to find the polynomial ζ as $\zeta = 272\sigma + 174$, satisfying the property $D(\sigma^2 + 10\sigma + 6)$.

3.1.5 Case 5

Consider the $\{\xi, \zeta\}$ where $\xi = 105\sigma + 67$ and $\zeta = 272\sigma + 174$ satisfying $D(\sigma^2 + 10\sigma + 6)$. To search for its extendability, let ϕ be any other polynomial with the same property. This can be written as $\xi\phi + \sigma^2 + 10\sigma + 6 = \rho^2$ and $\zeta\phi + \sigma^2 + 10\sigma + 6 = \eta^2$.

Applying the linear transformation, $\rho = \mu + (105\sigma + 67)\nu$ and $\eta = \mu + (272\sigma + 174)\nu$ and eliminating ϕ , we get the Pellian equation $\mu^2 - (28560\sigma^2 + 36494\sigma + 11658)\nu^2 = \sigma^2 + 10\sigma + 6$, with initial solution $\mu_0 = 169\sigma + 108, \nu_0 = 1$. The values of μ_0 and ν_0 can be employed to find the polynomial ϕ as $\phi = 715\sigma + 457$, satisfying the property $D(\sigma^2 + 10\sigma + 6)$.

3.2 Choice 2

For the different pairs of linear polynomials, we investigate for its extendability as a triple satisfying the property $D(10\omega^2 + 26\omega + 13)$.

3.2.1 Case 1

Consider the $\{\alpha, \beta\}$ where $\alpha = 3\omega + 1$ and $\beta = 5\omega + 3$ satisfying $D(10\omega^2 + 26\omega + 13)$. To search for its extendability, let γ be any other polynomial with the same property. This can be written as $\alpha\gamma + 10\omega^2 + 26\omega + 13 = \rho^2$ and $\beta\gamma + 10\omega^2 + 26\omega + 13 = \eta^2$.

Applying the linear transformation, $\rho = \mu + (3\omega + 1)\nu$ and $\eta = \mu + (5\omega + 3)\nu$ and eliminating γ , we get the Pellian equation $\mu^2 - (15\omega^2 + 14\omega + 3)\nu^2 = 10\omega^2 + 26\omega + 13$, with initial solution $\mu_0 = 5\omega + 4, \nu_0 = 1$. The values of μ_0 and ν_0 can be employed to find the polynomial γ as $\gamma = 18\omega + 12$, satisfying the property $D(10\omega^2 + 26\omega + 13)$.

3.2.2 Case 2

Consider the $\{\beta, \gamma\}$ where $\beta = 5\omega + 3$ and $\gamma = 18\omega + 12$ satisfying $D(10\omega^2 + 26\omega + 13)$. To search for its extendability, let δ be any other polynomial with the same property. This can be written as $\beta\delta + 10\omega^2 + 26\omega + 13 = \rho^2$ and $\gamma\delta + 10\omega^2 + 26\omega + 13 = \eta^2$.

Applying the linear transformation, $\rho = \mu + (5\omega + 3)\nu$ and $\eta = \mu + (18\omega + 12)\nu$ and eliminating δ , we get the Pellian equation $\mu^2 - (90\omega^2 + 114\omega + 36)\nu^2 = 10\omega^2 + 26\omega + 13$, with initial solution $\mu_0 = 10\omega + 7, \nu_0 = 1$. The values of μ_0 and ν_0 can be employed to find the polynomial δ as $\delta = 43\omega + 29$, satisfying the property $D(10\omega^2 + 26\omega + 13)$.

3.2.3 Case 3

Consider the $\{\gamma, \delta\}$ where $\gamma = 18\omega + 12$ and $\delta = 43\omega + 29$ satisfying $D(10\omega^2 + 26\omega + 13)$. To search for its extendability, let ξ be any other polynomial with the same property. This can be written as $\gamma\xi + 10\omega^2 + 26\omega + 13 = \rho^2$ and $\delta\xi + 10\omega^2 + 26\omega + 13 = \eta^2$.

Applying the linear transformation, $\rho = \mu + (18\omega + 12)\nu$ and $\eta = \mu + (43\omega + 29)\nu$ and eliminating ξ , we get the Pellian equation $\mu^2 - (774\omega^2 + 1038\omega + 348)\nu^2 = 10\omega^2 + 26\omega + 13$, with initial solution $\mu_0 = 28\omega + 19, \nu_0 = 1$. The values of μ_0 and ν_0 can be employed to find the polynomial ξ as $\xi = 117\omega + 79$, satisfying the property $D(10\omega^2 + 26\omega + 13)$.

3.2.4 Case 4

Consider the $\{\delta, \xi\}$ where $\delta = 43\omega + 29$ and $\xi = 117\omega + 79$ satisfying $D(10\omega^2 + 26\omega + 13)$. To search for its extendability, let ζ be any other polynomial with the same property. This can be written as $\delta\zeta + 10\omega^2 + 26\omega + 13 = \rho^2$ and $\xi\zeta + 10\omega^2 + 26\omega + 13 = \eta^2$.


Applying the linear transformation, $\rho = \mu + (43\omega + 29)\nu$ and $\eta = \mu + (117\omega + 79)\nu$ and eliminating ζ , we get the Pellian equation $\mu^2 - (5031\omega^2 + 6790\omega + 2291)\nu^2 = 10\omega^2 + 26\omega + 13$, with initial solution $\mu_0 = 71\omega + 48, \nu_0 = 1$. The values of μ_0 and ν_0 can be employed to find the polynomial ζ as $\zeta = 302\omega + 204$, satisfying the property $D(10\omega^2 + 26\omega + 13)$.

3.2.5 Case 5

Consider the $\{\xi, \zeta\}$ where $\xi = 117\omega + 79$ and $\zeta = 302\omega + 204$ satisfying $D(10\omega^2 + 26\omega + 13)$. To search for its extendability, let ϕ be any other polynomial with the same property. This can be written as $\xi\phi + 10\omega^2 + 26\omega + 13 = \rho^2$ and $\zeta\phi + 10\omega^2 + 26\omega + 13 = \eta^2$. Applying the linear transformation, $\rho = \mu + (117\omega + 79)\nu$ and $\eta = \mu + (302\omega + 204)\nu$ and eliminating ϕ , we get the Pellian equation $\mu^2 - (35334\omega^2 + 47726\omega + 16116)\nu^2 = 10\omega^2 + 26\omega + 13$, with initial solution $\mu_0 = 188\omega + 127, \nu_0 = 1$. The values of μ_0 and ν_0 can be employed to find the polynomial ϕ as $\phi = 795\omega + 537$, satisfying the property $D(10\omega^2 + 26\omega + 13)$.

4 CONCLUSION

All the triples considered in this paper cannot be extended to a quadruple for the particular choices of quadratic polynomial.



References

- [1] I. NIVEN, H.S. ZUCKERMAN, H. L. MONTGOMETRY, An Introduction to the Theory of Numbers, *Fifth Edition*, John Wiley and Sons, Inc., New York(1991).
www.fuchs-braun.com/media/532896481f9c1c47ffff8077ffff0.pdf
- [2] BO HE AND A. TOGBE, On the family of Diophantine triples $\{k + 1, 4k, 9k + 3\}$, *Period Math. Hunger*, 58(2009), 59 - 70.
<https://link.springer.com/article/10.1007/s10998-009-9059-6>
- [3] MANJU SOMANATH AND J. KANNAN, On A Class of Solutions for a Diophantine equation of Second degree. *International Journal of Pure and Applied Mathematics*, Special Issue, Vol. 117, No. 12(2017) 55-62.
<http://acadpubl.eu/jsi/2017-117-11-14/articles/12/7.pdf>
- [4] MANJU SOMANATH, J. KANNAN, K. RAJA AND V. SANGEETHA, On the Integer Solutions of the Pell's equation $x^2 = 17y^2 - 19^t$, *JP Journal of Applied Mathematics*, Vol. 15, Issue 2 (2017) 81-88.
www.iphsci.com/on_the_integer_solutions_abstract_references.pdf

